arXiv:1509.09215v1 [math.FA] 30 Sep 2015

VECTOR-VALUED CHARACTERS ON VECTOR-VALUED FUNCTION ALGEBRAS

MORTAZA ABTAHI

Abstract. Let A be a commutative Banach algebra and X be a compact space. The class of Banach A-valued function algebras on X consists of subalgebras of $C(X, A)$ with certain properties. We introduce the notion of A-characters on an A-valued function algebra $\mathscr{A}$ as homomorphisms from $\mathscr{A}$ into A that basically have the same properties as the evaluation homomorphisms $\mathcal{E}_x : f \mapsto f(x)$, with $x \in X$. For the so-called natural A-valued function algebras, such as $C(X, A)$ and $\mathrm{Lip}(X, A)$, we show that $\mathcal{E}_x$ $(x \in X)$ are the only A-characters. Vector-valued characters are utilized to identify vector-valued spectrums. When $A = \mathbb{C}$, Banach A-valued function algebras reduce to Banach function algebras, and A-characters reduce to characters.

1. Introduction and Preliminaries

In this paper, we consider commutative unital Banach algebras over the complex field $\mathbb{C}$. For more information on the theory of commutative Banach algebras, see the texts [3, 5, 14, 20].

1.1. **Characters.** Let A be a commutative Banach algebra with identity $\mathbf{1}$. Every nonzero homomorphism $\phi : A \to \mathbb{C}$ is called a *character* of A. The set of all characters of A is denoted by $\mathfrak{M}(A)$. It is well-known that $\mathfrak{M}(A)$ is nonempty and its elements are automatically continuous [14, Lemma 2.1.5]. For every $a \in A$, define $\hat{a} : \mathfrak{M}(A) \to \mathbb{C}$ by $\hat{a}(\phi) = \phi(a)$. The *Gelfand topology* of $\mathfrak{M}(A)$ is then the weakest topology on $\mathfrak{M}(A)$ that makes every $\hat{a}$ a continuous function. Equipped with the Gelfand topology, $\mathfrak{M}(A)$ is a compact Hausdorff space. The set $\hat{A} = \{\hat{a} : a \in A\}$ is a subalgebra of $C(\mathfrak{M}(A))$ and the *Gelfand transform* $A \to \hat{A}$, $a \mapsto \hat{a}$, is an algebra homomorphisms with $\operatorname{rad} A = \{a : \hat{a} = 0\}$ as its kernel. The algebra A is called *semisimple* if $\operatorname{rad} A = \mathbf{0}$.

1.2. **Complex Function Algebras.** Let X be a compact Hausdorff space and let $C(X)$ be the algebra of all continuous functions $f : X \to \mathbb{C}$ equipped with

Date: Received: xxxxxx; Revised: yyyyyy; Accepted: zzzzzz.

2010 *Mathematics Subject Classification.* Primary 46J10, 46H10; Secondary 46J20, 46E40.

Key words and phrases. Algebras of continuous vector-valued functions, Banach function algebras, Vector-valued function algebras, Characters, Maximal ideals.

the uniform norm $\|f\|_X = \sup\{|f(x)| : x \in X\}$. We recall that a subalgebra $\mathfrak{A}$ of $C(X)$ that separates the points of X and contains the constant functions is a *function algebra* on X. A function algebra $\mathfrak{A}$ equipped with some algebra norm $\|\cdot\|$ such that $\|f\|_X \leq \|f\|$, for all $f \in \mathfrak{A}$, is a *normed function algebra*. A complete normed function algebra is a *Banach function algebra*. If the norm of a Banach function algebra $\mathfrak{A}$ is equivalent to the uniform norm $\|\cdot\|_X$, then $\mathfrak{A}$ is a *uniform algebra*.

Let $\mathfrak{A}$ be a Banach function algebra on X. For every $x \in X$, the mapping $\varepsilon_x : \mathfrak{A} \to \mathbb{C}$, $\varepsilon_x(f) = f(x)$, is a character of $\mathfrak{A}$, and the mapping $J : X \to \mathfrak{M}(\mathfrak{A})$, $x \mapsto \varepsilon_x$, imbeds X homeomorphically as a compact subset of $\mathfrak{M}(\mathfrak{A})$. When J is surjective, one calls $\mathfrak{A}$ *natural* [5, Chapter 4]. For instance, $C(X)$ is a natural uniform algebra. We remark that every semisimple commutative unital Banach algebra A can be seen (through its Gelfand transform) as a natural Banach function algebra on $X = \mathfrak{M}(A)$. For more on function algebras see, for example, [7, 15].

1.3. **Vector-valued Function Algebras.** Let $(A, \|\cdot\|)$ be a commutative unital Banach algebra. The set of all A-valued continuous functions on X is denoted by $C(X, A)$. Note that $C(X) = C(X, \mathbb{C})$. For $f, g \in C(X, A)$ and $\alpha \in \mathbb{C}$, the pointwise operations $\alpha f + g$ and fg are defined in the obvious way. The *uniform norm* of each function $f \in C(X, A)$ is defined by $\|f\|_X = \sup\{\|f(x)\| : x \in X\}$. In this setting, $\big(C(X, A), \|\cdot\|_X\big)$ is a commutative unital Banach algebra.

Starting by Yood [19], in 1950, the character space of $C(X, A)$ has been studied by many authors. In 1957, Hausner [8] proved that τ is a character of $C(X, A)$ if, and only if, there exist a point $x \in X$ and a character $\phi \in \mathfrak{M}(A)$ such that $\tau(f) = \phi(f(x))$, for all $f \in C(X, A)$, whence $\mathfrak{M}(C(X, A))$ is homeomorphic to $X \times \mathfrak{M}(A)$. Recently, in [1], other characterizations of maximal ideals in $C(X, A)$ have been presented.

Analogous with complex function algebras, vector-valued function algebras are defined (Definition 2.1).

1.4. **Vector-valued Characters.** Let $\mathscr{A}$ be an A-valued function algebra on X. For each $x \in X$, consider the evaluation homomorphism $\mathcal{E}_x : \mathscr{A} \to A$, $x \mapsto f(x)$. These A-valued homomorphisms are included in a certain class of homomorphisms that will be introduced and studied in Section 3 under the name of *vector-valued characters*. More precisely, a homomorphism $\Psi : \mathscr{A} \to A$ is called an *A-character* if $\Psi(\mathbf{1}) = \mathbf{1}$ and $\phi(\Psi f) = \Psi(\phi \circ f)$, for all $f \in \mathscr{A}$ and $\phi \in \mathfrak{M}(A)$. See Definition 3.1. The set of all A-characters of $\mathscr{A}$ will be denoted by $\mathfrak{M}_A(\mathscr{A})$. Note that when $A = \mathbb{C}$, we have $\mathfrak{M}_A(\mathscr{A}) = \mathfrak{M}_{\mathbb{C}}(\mathscr{A}) = \mathfrak{M}(\mathscr{A})$.

An application of vector-valued characters are presented in the forthcoming paper [2] to identify the vector-valued spectrum of functions $f \in \mathscr{A}$. It is known that the spectrum of an element $a \in A$ is equal to $\operatorname{sp}(a) = \{\phi(a) : \phi \in \mathfrak{M}(A)\}$. In [2] the A-valued spectrum $\vec{\operatorname{sp}}_A(f)$ of functions $f \in \mathscr{A}$ are studied and it is proved that, under certain conditions,

$$\vec{\operatorname{sp}}_A(f) = \{\Psi(f) : \Psi \in \mathfrak{M}_A(\mathscr{A})\}.$$

1.5. **Notations and conventions.** Since, in this paper, we are dealing with different types of functions and algebras, ambiguity may arise. Hence, a clear declaration of notations and conventions is given here.

(1) Throughout the paper, X is a compact Hausdorff space, and A is a commutative *semisimple* Banach algebra with identity $\mathbf{1}$. The set of invertible elements of A is denoted by $\operatorname{Inv}(A)$.
(2) If $f : X \to \mathbb{C}$ is a function and $a \in A$, we write fa to denote the A-valued function $X \to A$, $x \mapsto f(x)a$. If $\mathfrak{A}$ is a complex-valued function algebra on X, we let $\mathfrak{A}A$ be the linear span of $\{fa : f \in \mathfrak{A},\, a \in A\}$. Hence, an element $f \in \mathfrak{A}A$ is of the form $f = f_1a_1 + \cdots + f_na_n$ with $f_j \in \mathfrak{A}$ and $a_j \in A$.
(3) Given an element $a \in A$, we use the same notation a for the constant function $X \to A$ given by $a(x) = a$, for all $x \in X$, and consider A as a closed subalgebra of $C(X, A)$. Since we assume that A has an identity $\mathbf{1}$, we identify $\mathbb{C}$ with the closed subalgebra $\mathbb{C}\mathbf{1}$ of A, and thus every continuous function $f : X \to \mathbb{C}$ can be seen as the continuous A-valued function $X \to A$, $x \mapsto f(x)\mathbf{1}$; we use the same notation f for this A-valued function, and regard $C(X)$ as a closed subalgebra of $C(X, A)$.
(4) To every continuous function $f : X \to A$, we correspond the function

$$\tilde{f} : \mathfrak{M}(A) \to C(X), \quad \tilde{f}(\phi) = \phi \circ f.$$

If $\mathcal{I}$ is a family of continuous A-valued functions on X, we define

$$\phi[\mathcal{I}] = \{\phi \circ f : f \in \mathcal{I}\} = \{\tilde{f}(\phi) : f \in \mathcal{I}\}.$$

2. Vector-valued Function Algebras

In this section, we introduce the notion of vector-valued function algebras as subalgebras of $C(X, A)$ with certain properties. Vector-valued function algebras have been defined and studied by others (e.g. [16]).

Definition 2.1. Let X be a compact Hausdorff space, and $(A, \|\cdot\|)$ be a commutative unital Banach algebra. An *A-valued function algebra* on X is a subalgebra $\mathscr{A}$ of $C(X, A)$ such that (1) $\mathscr{A}$ contains all the constant functions $X \to A$, $x \mapsto a$, with $a \in A$, and (2) $\mathscr{A}$ separates the points of X in the sense that, for every pair

x, y of distinct points in X, there exists $f \in A$ such that $f(x) \neq f(y)$. A *normed A-valued function algebra* on X is an A-valued function algebra $\mathscr{A}$ on X endowed with some algebra norm $|||\cdot|||$ such that the restriction of $|||\cdot|||$ to A is equivalent to the original norm $\|\cdot\|$ of A, and $\|f\|_X \leq |||f|||$, for every $f \in \mathscr{A}$. A complete normed A-valued function algebra is called a *Banach A-valued function algebra*. A Banach A-valued function algebra $\mathscr{A}$ is called an *A-valued uniform algebra* if the given norm of $\mathscr{A}$ is equivalent to the uniform norm $\|\cdot\|_X$.

If there is no risk of confusion, instead of $|||\cdot|||$, we use the same notation $\|\cdot\|$ for the norm of $\mathscr{A}$.

Let $\mathscr{A}$ be an A-valued function algebra on X. For every $x \in X$, define $\mathcal{E}_x : \mathscr{A} \to A$ by $\mathcal{E}_x(f) = f(x)$. We call $\mathcal{E}_x$ the *evaluation homomorphism* at x. Our definition of Banach A-valued function algebras implies that every evaluation homomorphism $\mathcal{E}_x$ is continuous. As it is mentioned in [16], if the condition $\|f\|_X \leq |||f|||$, for all $f \in \mathscr{A}$, is replaced by the requirement that every evaluation homomorphism $\mathcal{E}_x$ is continuous, then one can find some constant M such that

$$\|f\|_X \leq M|||f||| \quad (f \in \mathscr{A}).$$

2.1. **Admissible algebras.** Given a complex-valued function algebra $\mathfrak{A}$ and an A-valued function algebra $\mathscr{A}$ on X, according to [16, Definition 2.1], the quadruple $(X, A, \mathfrak{A}, \mathscr{A})$ is *admissible* if $\mathfrak{A}$ is natural, $\mathfrak{A}A \subset \mathscr{A}$, and

$$\{\phi \circ f : \phi \in \mathfrak{M}(A), f \in \mathscr{A}\} \subset \mathfrak{A}.$$

Taking this into account, we make the following definition.

Definition 2.2. An A-valued function algebra $\mathscr{A}$ is said to be *admissible* if

$$\{\phi \circ f : \phi \in \mathfrak{M}(A), f \in \mathscr{A}\} \subset \mathscr{A}. \tag{2.1}$$

When $\mathscr{A}$ is admissible, we set $\mathfrak{A} = C(X) \cap \mathscr{A}$ which is the subalgebra of $\mathscr{A}$ consisting of all complex-valued functions in $\mathscr{A}$ and forms a complex-valued function algebra by itself. Note that, in this case, $\mathfrak{A} = \phi[\mathscr{A}]$, for all $\phi \in \mathfrak{M}(A)$.

Example 2.3. Let $\mathfrak{A}$ be a complex-valued function algebra on X. Then $\mathfrak{A}A$ is an admissible A-valued function algebra on X. Hence, the uniform closure of $\mathfrak{A}A$ in $C(X, A)$ is an admissible A-valued uniform algebra (Proposition 2.5 below).

Another example of an admissible Banach A-valued function algebra is the A-valued Lipschitz function algebra $\mathrm{Lip}(X, A)$, where X is a compact metric space. Recently, in [6], the character space and Śilov boundary of $\mathrm{Lip}(X, A)$ has been studied. For certain properties of vector-valued Lipschitz spaces, see [4] and [12, 13].

The following shows that not all vector-valued function algebras are admissible.

Example 2.4. Let K be a compact subset of $\mathbb{C}$ which is not polynomially convex so that $P(K) \neq R(K)$. For example, let $K = \mathbb{T}$ be the unit circle. Set

$$\mathscr{A} = \{(f_p, f_r) : f_p \in P(K), f_r \in R(K)\}.$$

Then $\mathscr{A}$ is a uniformly closed subalgebra of $C(K, \mathbb{C}^2)$ which contains all the constant functions $(\alpha, \beta) \in \mathbb{C}^2$ and separates the points of K. Hence $\mathscr{A}$ is a $\mathbb{C}^2$-valued uniform algebra on K. Let $\mathbf{1} = (1, 1)$ be the unit element of $\mathbb{C}^2$, and let π_1 and π_2 be the coordinate projections of $\mathbb{C}^2$. Then $\mathfrak{M}(\mathbb{C}^2) = \{\pi_1, \pi_2\}$, and

$$\pi_1[\mathscr{A}]\mathbf{1} = \{f\mathbf{1} : f \in P(K)\} = \{(f, f) : f \in P(K)\},$$
$$\pi_2[\mathscr{A}]\mathbf{1} = \{f\mathbf{1} : f \in R(K)\} = \{(f, f) : f \in R(K)\}.$$

We see that $\pi_1[\mathscr{A}] \subset \mathscr{A}$ while $\pi_2[\mathscr{A}] \not\subset \mathscr{A}$. Hence $\mathscr{A}$ is not admissible.

Proposition 2.5. *Suppose $\mathscr{A}$ is an admissible A-valued function algebra on X. Then the uniform closure $\bar{\mathscr{A}}$ is an admissible A-valued uniform algebra on X and $\bar{\mathfrak{A}} = C(X) \cap \bar{\mathscr{A}}$.*

Proof. Recall that $\mathfrak{A} = C(X) \cap \mathscr{A}$. The fact that $\bar{\mathscr{A}}$ is an A-valued uniform algebra is clear. The inclusion $\bar{\mathfrak{A}} \subset C(X) \cap \bar{\mathscr{A}}$ is also obvious. Take $f \in C(X) \cap \bar{\mathscr{A}}$. Then, there exists a sequence $\{f_n\}$ of A-valued functions in $\mathscr{A}$ such that $f_n \to f$ uniformly on X. For some $\phi \in \mathfrak{M}(A)$, take $g_n = \phi \circ f_n$. Then $g_n \in \mathfrak{A}$ and, since $f = \phi \circ f$, we have

$$\|g_n - f\|_X = \|\phi \circ f_n - \phi \circ f\|_X \leq \|f_n - f\|_X \to 0.$$

Therefore, $g_n \to f$ uniformly on X and thus $f \in \bar{\mathfrak{A}}$. □

2.2. **Certain vector-valued uniform algebras.** Let $\mathfrak{A}_0$ be a complex function algebra on X, and let $\mathfrak{A}$ and $\mathscr{A}$ be the uniform closures of $\mathfrak{A}_0$ and $\mathfrak{A}_0 A$, in $C(X)$ and $C(X, A)$, respectively. Then $\mathscr{A}$ is an admissible A-valued uniform algebra on X with $\mathfrak{A} = C(X) \cap \mathscr{A}$. The algebra $\mathscr{A}$ is isometrically isomorphic to the injective tensor product $\mathfrak{A} \hat{\otimes}_\epsilon A$ (cf. [14, Proposition 1.5.6]). To see this, let $T : \mathfrak{A} \otimes A \to \mathscr{A}$ be the unique linear mapping, given by [3, Theorem 42.6], such that $T(f \otimes a)(x) = f(x)a$, for all $x \in X$. Let $\mathfrak{A}_1^*$ and A_1^* denote the closed unit ball of $\mathfrak{A}^*$ and A^*, respectively, and let $\|\cdot\|_\epsilon$ denote the injective tensor norm.

Then

$$\begin{aligned}\Big\|T\Big(\sum_{i=1}^n f_i\otimes a_i\Big)\Big\|_X &= \sup_{x\in X}\Big\|\sum_{i=1}^n f_i(x)a_i\Big\| = \sup_{x\in X}\sup_{\nu\in A_1^*}\Big|\sum_{i=1}^n f_i(x)\nu(a_i)\Big| \\ &= \sup_{\nu\in A_1^*}\Big\|\sum_{i=1}^n f_i(\cdot)\nu(a_i)\Big\|_X = \sup_{\nu\in A_1^*}\sup_{\mu\in\mathfrak{A}_1^*}\Big|\mu\Big(\sum_{i=1}^n f_i(\cdot)\nu(a_i)\Big)\Big| \\ &= \sup_{\mu\in\mathfrak{A}_1^*}\sup_{\nu\in A_1^*}\Big|\sum_{i=1}^n \mu(f_i)\nu(a_i)\Big| = \Big\|\sum_{i=1}^n f_i\otimes a_i\Big\|_\epsilon.\end{aligned}$$

Hence T extends to an isometry $\overline{T}$ from $\mathfrak{A}\,\hat{\otimes}_\epsilon A$ into $\mathscr{A}$. Since the range of T contains $\mathfrak{A}_0 A$, which is dense in $\mathscr{A}$, the range of $\overline{T}$ is the whole of $\mathscr{A}$. We remark that, by a theorem of Tomiyama [18], the maximal ideal space $\mathfrak{M}(\mathscr{A})$ of $\mathscr{A}$ is homeomorphic to $\mathfrak{M}(\mathfrak{A})\times\mathfrak{M}(A)$.

Now, let K be a compact subset of $\mathbb{C}$. Associated with K there are three vector-valued uniform algebras in which we are interested. Let $P_0(K,A)$ be the algebra of the restriction to K of all polynomials $p(z)=a_nz^n+\cdots+a_1z+a_0$ with coefficients $a_0,a_1,\ldots,a_n$ in A. Let $R_0(K,A)$ be the algebra of the restriction to K of all rational functions of the form $p(z)/q(z)$, where $p(z)$ and $q(z)$ are polynomials with coefficients in A, and $q(\lambda)\in\mathrm{Inv}(A)$, whenever $\lambda\in K$. And let $H_0(K,A)$ be the algebra of all A-valued functions on K having a holomorphic extension to a neighbourhood of K.

When $A=\mathbb{C}$, we drop A and write $P_0(K)$, $R_0(K)$ and $H_0(K)$. Their uniform closures in $C(K)$, denoted by $P(K)$, $R(K)$ and $H(K)$, are complex uniform algebras (for more on complex uniform algebras, see [7] or [15]).

The algebras $P_0(K,A)$, $R_0(K,A)$ and $H_0(K,A)$ are A-valued function algebras on K, and their uniform closures in $C(K,A)$, denoted by $P(K,A)$, $R(K,A)$ and $H(K,A)$, are A-valued uniform algebras. It obvious that

$$P(K,A)\subset R(K,A)\subset H(K,A).$$

Every polynomial $p(z)=a_nz^n+\cdots+a_1z+a_0$ in $P_0(K,A)$ can be written as $p(z)=p_0(z)a_0+p_1(z)a_1+\cdots+p_n(z)a_n$, where $p_0,p_1,\ldots,p_n$ are polynomials in $P_0(K)$. Thus $P_0(K,A)=P_0(K)A$. The above discussion shows that $P(K,A)$ is isometrically isomorphic to $P(K)\,\hat{\otimes}_\varepsilon A$. The maximal ideal space of $P(K)$ is homeomorphic to $\hat{K}$, the polynomially convex hull of K (e.g. [15, Section 5.2]). The maximal ideal space of $P(K,A)$ is, therefore, homeomorphic to $\hat{K}\times\mathfrak{M}(A)$; see also [16].

Runge's classical approximation theorem states that if Λ is a subset of $\mathbb{C}$ such that Λ has nonempty intersection with each bounded component of $\mathbb{C}\setminus K$, then

every function $f \in H_0(K)$ can be approximated uniformly on K by rational functions with poles only among the points of Λ and at infinity ([3, Theorem 7.7]). In particular, $R(K) = H(K)$. The following is a version of Runge's theorem for vector-valued functions.

Theorem 2.6 (Runge)**.** *Let K be a compact subset of $\mathbb{C}$, and let Λ be a subset of $\mathbb{C}\setminus K$ having nonempty intersection with each bounded component of $\mathbb{C}\setminus K$. Then every function $f \in H_0(K, A)$ can be approximated uniformly on K by A-valued rational functions of the form*

$$r(z) = r_1(z)a_1 + r_2(z)a_2 + \cdots + r_n(z)a_n, \tag{2.2}$$

where $r_i(z)$, for $1 \le i \le n$, are rational functions in $R_0(K)$ with poles only among the points of Λ and at infinity, and $a_1, a_2, \ldots, a_n \in A$.

Proof. Take $f \in H_0(K, A)$. Then, there exists an open set D such that $K \subset D$ and $f : D \to A$ is holomorphic. We use the same notation as in [3]. We let E be a punched disc envelope for (K, D); see [3, Definition 6.2]. The Cauchy theorem and the Cauchy integral formula are also valid for Banach space valued holomorphic functions (see the remark after Corollary 6.6 in [3] and [17, Theorem 3.31]). Therefore,

$$f(z) = \frac{1}{2\pi \mathrm{i}} \int_{\partial E} \frac{f(s)}{s - z} ds \quad (z \in K).$$

Then, by [3, Proposition 6.5], one can write

$$f(z) = \sum_{n=0}^{\infty} \alpha_n (z - z_0)^n + \sum_{j=1}^{m} \sum_{n=1}^{\infty} \frac{\beta_{jn}}{(z - z_j)^n} \quad (z \in K), \tag{2.3}$$

where $z_0 \in \mathbb{C}$, $z_1, z_2, \ldots, z_m \in \mathbb{C} \setminus K$, and the coefficients α_n, β_{jn} belong to A. Note that the series in (2.3) converges uniformly on K.

So far, we have seen that f can be approximated uniformly on K by A-valued rational functions of the form (2.2), where $r_i(z)$, for $1 \le i \le n$, are rational functions in $R_0(K)$ with poles just outside K. Using Runge's classical theorem, each $r_i(z)$ can be approximated uniformly on K by rational functions with poles only among the points of Λ and at infinity. Hence, we conclude that f can be approximated uniformly on K by rational functions of the form (2.2) with preassigned poles. □

As a consequence of the above theorem, we see that the uniform closures of $R_0(K)A$, $H_0(K)A$, $R_0(K, A)$ and $H_0(K, A)$ are all the same. In particular,

$$R(K, A) = H(K, A).$$

Corollary 2.7. *The algebra $R(K,A)$ is isometrically isomorphic to $R(K)\hat{\otimes}_\varepsilon A$ and, therefore, $\mathfrak{M}(R(K,A))$ is homeomorphic to $K\times\mathfrak{M}(A)$.*

We remark that the equality $\mathfrak{M}(R(K,A))=K\times\mathfrak{M}(A)$ is proved in [16]. The authors, however, did not notice the equality $R(K,A)=R(K)\hat{\otimes}_\varepsilon A$.

3. Vector-valued Characters

Let $\mathscr{A}$ be a Banach A-valued function algebra on X, and consider the point evaluation homomorphisms $\mathcal{E}_x:\mathscr{A}\to A$. These kind of homomorphisms enjoy the following properties:

- $\mathcal{E}_x(a)=a$, for all $a\in A$,
- $\mathcal{E}_x(\phi\circ f)=\phi(\mathcal{E}_x f)$, for all $f\in\mathscr{A}$ and $\phi\in\mathfrak{M}(A)$,
- If $\mathscr{A}$ is admissible (with $\mathfrak{A}=C(X)\cap\mathscr{A}$) then $\mathcal{E}_x|_{\mathfrak{A}}$ is a character of $\mathfrak{A}$, namely, the evaluation character ε_x.

We now introduce the class of all homomorphisms from $\mathscr{A}$ into A having the same properties as the point evaluation homomorphisms $\mathcal{E}_x$ $(x\in X)$.

Definition 3.1. Let $\mathscr{A}$ be an admissible A-valued function algebra on X. A homomorphism $\Psi:\mathscr{A}\to A$ is called an *A-character* if $\Psi(\mathbf{1})=\mathbf{1}$ and $\phi(\Psi f)=\Psi(\phi\circ f)$, for all $f\in\mathscr{A}$ and $\phi\in\mathfrak{M}(A)$. The set of all A-characters of $\mathscr{A}$ is denoted by $\mathfrak{M}_A(\mathscr{A})$.

That every A-character $\Psi:\mathscr{A}\to A$ satisfies $\Psi(a)=a$, for all $a\in A$, is easy to see. In fact, since $\phi(\Psi(a))=\Psi(\phi(a))=\phi(a)$, for all $\phi\in\mathfrak{M}(A)$, and A is semisimple, we get $\Psi(a)=a$.

Proposition 3.2. *Let $\Psi:\mathscr{A}\to A$ be a linear operator such that $\Psi(\mathbf{1})=\mathbf{1}$ and $\phi(\Psi f)=\Psi(\phi\circ f)$, for all $f\in\mathscr{A}$ and $\phi\in\mathfrak{M}(A)$. Then, the following are equivalent:*

(i) *Ψ is an A-character,*
(ii) *$\Psi(f)\neq\mathbf{0}$, for every $f\in\mathrm{Inv}(\mathscr{A})$,*
(iii) *$\Psi(f)\neq\mathbf{0}$, for every $f\in\mathrm{Inv}(\mathfrak{A})$,*
(iv) *if $\psi=\Psi|_{\mathfrak{A}}$, then $\psi\in\mathfrak{M}(\mathfrak{A})$.*

Proof. (i) $\Rightarrow$ (ii) $\Rightarrow$ (iii) is clear. The implication (iii) $\Rightarrow$ (iv) follows from [17, Theorem 10.9]. To prove (iv) $\Rightarrow$ (i), take $f,g\in\mathscr{A}$. For every $\phi\in\mathfrak{M}(A)$, we have

$$\phi(\Psi(fg))=\Psi(\phi\circ fg)=\psi((\phi\circ f)(\phi\circ g))=\psi(\phi\circ f)\psi(\phi\circ g)=\phi(\Psi(f)\Psi(g)).$$

Since A is semisimple, we get $\Psi(fg)=\Psi(f)\Psi(g)$. □

Every A-character is automatically continuous by Johnson's theorem [11]. If A is a uniform algebra and $\mathscr{A}$ is an A-valued uniform algebra, we even have $\|\Psi\| = 1$, for all A-character Ψ.

Proposition 3.3. *Let Ψ_1 and Ψ_2 be A-characters on $\mathscr{A}$, and set $\psi_1 = \Psi_1|_{\mathfrak{A}}$ and $\psi_2 = \Psi_2|_{\mathfrak{A}}$. The following are equivalent:*

(i) $\Psi_1 = \Psi_2$,
(ii) $\ker \Psi_1 = \ker \Psi_2$,
(iii) $\ker \psi_1 = \ker \psi_2$,
(iv) $\psi_1 = \psi_2$.

Proof. The implication (i) $\Rightarrow$ (ii) is obvious. The implication (ii) $\Rightarrow$ (iii), follows from the fact that $\ker \psi_i = \ker \Psi_i \cap \mathfrak{A}$, for $i = 1, 2$. The implication (iii) $\Rightarrow$ (iv) follows form [17, Theorem 11.5]. Finally, if we have (iv), then, for every $f \in \mathscr{A}$,

$$\phi(\Psi_1(f)) = \psi_1(\phi(f)) = \psi_2(\phi(f)) = \phi(\Psi_2(f)) \quad (\phi \in \mathfrak{M}(A)).$$

Since A is semisimple, we get $\Psi_1(f) = \Psi_2(f)$, for all $f \in \mathscr{A}$. □

Definition 3.4. Let $\mathscr{A}$ be an admissible A-valued function algebra on X. Given a character $\psi \in \mathfrak{M}(\mathfrak{A})$, if there exists an A-character Ψ on $\mathscr{A}$ such that $\Psi|_{\mathfrak{A}} = \psi$, then we say that ψ *lifts* to the A-character Ψ.

Proposition 3.3 shows that, if $\psi \in \mathfrak{M}(\mathfrak{A})$ lifts to Ψ_1 and Ψ_2, then $\Psi_1 = \Psi_2$. For every $x \in X$, the unique A-character to which the evaluation character ε_x lifts is the evaluation homomorphism $\mathcal{E}_x$. In the following, we investigate conditions under which every character $\psi \in \mathfrak{M}(\mathfrak{A})$ lifts to some A-character Ψ. To proceed, we need some definitions, notations and auxiliary results.

Suppose $\mathscr{A}$ is an admissible Banach A-valued function algebra on X with $\mathfrak{A} = C(X) \cap \mathscr{A}$. Then $\mathfrak{A}$ is a Banach function algebra. Since Banach function algebras are semisimple, by Johnson's theorem [11], for every $\phi \in \mathfrak{M}(A)$, the composition operator $C_\phi : \mathscr{A} \to \mathfrak{A}$, defined by $C_\phi(f) = \phi \circ f$, is a continuous homomorphism. Hence, for some constant M_ϕ, we have

$$\|\phi \circ f\| \leq M_\phi \|f\| \quad (f \in \mathscr{A}). \tag{3.1}$$

Now, for every $f \in \mathscr{A}$ consider the function $\tilde{f} : \mathfrak{M}(A) \to \mathfrak{A}$, $\phi \mapsto \phi \circ f$. Set $\mathcal{X} = \mathfrak{M}(A)$ and $\tilde{\mathscr{A}} = \{\tilde{f} : f \in \mathscr{A}\}$. Suppose every $\tilde{f}$ is continuous, that is, if $\phi_\alpha \to \phi$ in the Gelfand topology of $\mathfrak{M}(A)$, then $\phi_\alpha \circ f \to \phi \circ f$ in $\mathfrak{A}$ (this is the case for uniform algebras; see Corollary 3.6). Then $\tilde{\mathscr{A}}$ is an $\mathfrak{A}$-valued function algebra on $\mathcal{X}$. In Theorem 3.9, we will discuss conditions under which $\tilde{\mathscr{A}}$ is admissible. Since every composite operator $C_\phi : \mathscr{A} \to \mathfrak{A}$, $f \mapsto \phi \circ f$, with $\phi \in \mathfrak{M}(A)$, is

continuous, we see that the inclusion map $\tilde{\mathscr{A}} \hookrightarrow \mathfrak{A}^{\mathcal{X}}$ is continuous, where $\mathfrak{A}^{\mathcal{X}}$ is given the cartesian product topology. The Closed Graph Theorem then implies that the inclusion map $\tilde{\mathscr{A}} \hookrightarrow C(\mathcal{X}, \mathfrak{A})$ is continuous so that, for some constant M, we have

$$\|\phi \circ f\| \leq M\|f\| \quad (\phi \in \mathfrak{M}(A),\ f \in \mathscr{A}). \tag{3.2}$$

Note that $\phi \circ f \in C(X)$ and $\|\phi \circ f\|_X \leq \|\phi\|\|f\|_X$, for every $\phi \in A^*$, $f \in \mathscr{A}$. We extend $\tilde{f}$ to a mapping from A^* to $C(X)$, and we still denoted this extension is by $\tilde{f}$.

Proposition 3.5. *With respect to the w*-topology of A^* and the uniform topology of $C(X)$, every mapping $\tilde{f} : A^* \to C(X)$ is continuous on bounded subsets of A^*.*

Proof. Let $\{\phi_\alpha\}$ be a net in A^* that converges, in the w*-topology, to some $\phi_0 \in A^*$ and suppose $\|\phi_\alpha\| \leq M$, for all α. Take $\varepsilon > 0$ and set, for every $x \in X$,

$$V_x = \{s \in X : \|f(s) - f(x)\| < \varepsilon\}.$$

Then $\{V_x : x \in X\}$ is an open covering of the compact space X. Hence, there exist finitely many points $x_1, \ldots, x_n$ in X such that $X \subset V_{x_1} \cup \cdots \cup V_{x_n}$. Set

$$U_0 = \big\{\phi \in A^* : \big|\phi\big(f(x_i)\big) - \phi_0\big(f(x_i)\big)\big| < \varepsilon,\ 1 \leq i \leq n\big\}.$$

The set U_0 is an open neighbourhood of ϕ_0 in the w*-topology. Since $\phi_\alpha \to \phi_0$, there exists α_0 such that $\phi_\alpha \in U_0$ for $\alpha \geq \alpha_0$. If $x \in X$, then $\|f(x) - f(x_i)\| < \varepsilon$, for some $i \in \{1, \ldots, n\}$, and thus, for $\alpha \geq \alpha_0$,

$$\begin{aligned} |\phi_\alpha \circ f(x) - \phi_0 \circ f(x)| &\leq |\phi_\alpha(f(x)) - \phi_\alpha(f(x_i))| + |\phi_\alpha(f(x_i)) - \phi_0(f(x_i))| \\ &\quad + |\phi_0(f(x_i)) - \phi_0(f(x))| \\ &< M\varepsilon + \varepsilon + \|\phi_0\|\varepsilon. \end{aligned}$$

Since $x \in X$ is arbitrary, we get $\|\phi_\alpha \circ f - \phi_0 \circ f\|_X \leq \varepsilon(M + \|\phi_0\| + 1)$, for $\alpha \geq \alpha_0$. □

Since $\mathfrak{M}(A)$ is a bounded subset of A^*, we get the following result for uniform algebras.

Corollary 3.6. *Let $\mathscr{A}$ be an admissible A-valued uniform algebra on A. Then $\tilde{f} \in C(\mathfrak{M}(A), \mathfrak{A})$, for every $f \in \mathscr{A}$, and $\tilde{\mathscr{A}} = \{\tilde{f} : f \in \mathscr{A}\}$ is an $\mathfrak{A}$-valued uniform algebra on $\mathfrak{M}(A)$.*

Example 3.7. Let K_1 and K_2 be compact subsets of the complex plane $\mathbb{C}$. Let $\mathfrak{A} = P(K_1)$ and $A = R(K_2)$. It is well-known that $\mathfrak{M}(R(K_2)) = K_2$. Set

$$\mathscr{A} = P(K_1, R(K_2)) = P(K_1) \,\hat{\otimes}_\varepsilon\, R(K_2).$$

Every $f \in P(K_1, R(K_2))$ induces a map $\tilde{f} : K_2 \to P(K_1)$ given by

$$\tilde{f}(z_2)(z_1) = f(z_1)(z_2).$$

We see that $\tilde{\mathscr{A}} = R(K_2, P(K_1)) = R(K_2) \hat{\otimes}_\varepsilon P(K_1)$.

Lemma 3.8. *For every $f \in \mathscr{A}$, if $\tilde{f} : \mathfrak{M}(A) \to \mathfrak{A}$ is scalar-valued, then f is a constant function and, therefore, $\tilde{f} = \hat{a}$, for some $a \in A$.*

Proof. Fix a point $x_0 \in A$ and let $a = f(x_0)$. The function $\tilde{f}$ being scalar-valued means that, for every $\phi \in \mathfrak{M}(A)$, there is a complex number λ such that $\tilde{f}(\phi) = \phi \circ f = \lambda$. This means that $\phi \circ f$ is a constant function on X so that

$$\phi(f(x)) = \phi(f(x_0)) = \phi(a) \quad (x \in X). \tag{3.3}$$

Since A is semisimple and (3.3) holds for every $\phi \in \mathfrak{M}(A)$, we must have $f(x) = a$, for all $x \in X$. Thus, $\tilde{f} = \hat{a}$. □

We are now ready to state and prove the main result of the section.

Theorem 3.9. *Let $\mathscr{A}$ be an admissible Banach A-valued function algebra on X, and let E be the linear span of $\mathfrak{M}(A)$ in A^*. The following statements are equivalent.*

(i) *for every $\psi \in \mathfrak{M}(\mathfrak{A})$ and $f \in \mathscr{A}$, the mapping $g : E \to \mathbb{C}$, defined by $g(\phi) = \psi(\phi \circ f)$, is continuous with respect to the w*-topology of E;*

(ii) *every $\psi \in \mathfrak{M}(\mathfrak{A})$ lifts to an A-character $\Psi : \mathscr{A} \to A$;*

(iii) *every $f \in \mathscr{A}$ has a unique extension $F : \mathfrak{M}(\mathfrak{A}) \to A$ such that*

$$\phi(F(\psi)) = \psi(\phi \circ f) \quad (\psi \in \mathfrak{M}(\mathfrak{A}),\ \phi \in \mathfrak{M}(A));$$

Moreover, if the functions $\tilde{f} : \mathfrak{M}(A) \to \mathfrak{A}$, where $f \in \mathscr{A}$, are all continuous so that $\tilde{\mathscr{A}} = \{\tilde{f} : f \in \mathscr{A}\}$ is an $\mathfrak{A}$-valued function algebra on $\mathfrak{M}(A)$, then the above statements are equivalent to

(iv) *$\tilde{\mathscr{A}}$ is admissible.*

Proof. (i) ⇒ (ii): Fix $\psi \in \mathfrak{M}(\mathfrak{A})$ and $f \in \mathscr{A}$, and consider the function $g : E \to \mathbb{C}$ defined by $g(\phi) = \psi(\phi \circ f)$. Since $\phi \circ f \in \mathfrak{A}$, for every $\phi \in E$, we see that g is a well-defined linear functional on E. Endowed with the w*-topology, A^* is a locally convex space with A as its dual. Since g is w*-continuous, by the Hahn-Banach extension theorem for locally convex spaces, [17, Theorem 3.6], there is a w*-continuous linear functional G on A^* that extends g. Hence $G = \hat{a}$, for some $a \in A$, and since A is semisimple, a is unique. Now, define $\Psi(f) = a$. Then

$$\phi(\Psi(f)) = \hat{a}(\phi) = g(\phi) = \psi(\phi \circ f) \quad (\phi \in \mathfrak{M}(A)).$$

It is easily seen that $\Psi : \mathscr{A} \to A$ is an A-character and $\Psi|_{\mathfrak{A}} = \psi$.

(ii) $\Rightarrow$ (iii): Fix $f \in \mathscr{A}$ and define $F : \mathfrak{M}(\mathfrak{A}) \to A$ by $F(\psi) = \Psi(f)$, where Ψ is the unique A-character of $\mathscr{A}$ to which ψ lifts. Considering the identification $x \mapsto \varepsilon_x$ and the fact that each ε_x lifts to $\mathcal{E}_x : f \mapsto f(x)$, we get

$$F(x) = F(\varepsilon_x) = \mathcal{E}_x(f) = f(x) \quad (x \in X).$$

So, $F|_X = f$. Also, $\phi(F(\psi)) = \phi(\Psi(f)) = \psi(\phi \circ f)$, for all $\phi \in \mathfrak{M}(A)$ and $\psi \in \mathfrak{M}(\mathfrak{A})$.

(iii) $\Rightarrow$ (i): Fix $\psi \in \mathfrak{M}(\mathfrak{A})$ and $f \in \mathscr{A}$, and put $a = F(\psi)$, where F is the unique extension of f to $\mathfrak{M}(\mathfrak{A})$ given by (iii). Then $g(\phi) = \hat{a}(\phi)$, for every $\phi \in E$, which is, obviously, a continuous function with respect to the w*-topology of E.

Finally, suppose $\tilde{\mathscr{A}} = \{\tilde{f} : f \in \mathscr{A}\}$ is an $\mathfrak{A}$-valued function algebra on $\mathfrak{M}(A)$. We prove (ii) $\Rightarrow$ (iv) $\Rightarrow$ (i). If $\mathscr{A}$ satisfies (ii), then

$$\psi \circ \tilde{f} = \widehat{\Psi(f)} \in \hat{A} \subset \tilde{\mathscr{A}}, \quad (f \in \mathscr{A}, \psi \in \mathfrak{M}(\mathfrak{A})).$$

This means that $\{\psi \circ \tilde{f} : \psi \in \mathfrak{M}(\mathfrak{A}), f \in \mathscr{A}\} \subset \tilde{\mathscr{A}}$ and thus $\tilde{\mathscr{A}}$ is admissible. Conversely, suppose $\tilde{\mathscr{A}}$ is admissible. Then, for every $\psi \in \mathfrak{M}(\mathfrak{A})$ and $f \in \mathscr{A}$, the function $\psi \circ \tilde{f}$ is a complex-valued function in $\tilde{\mathscr{A}}$. By Lemma 3.8, $\psi \circ \tilde{f}$ belongs to $\hat{A}$ so that $\psi \circ \tilde{f} = \hat{a}$, for some $a \in A$. Hence, $g(\phi) = \hat{a}(\phi)$, for every $\phi \in E$, and g is w*-continuous. □

In the following, we regard A and $\mathfrak{A}$ as subalgebras of $C(\mathfrak{M}(A))$ and $C(\mathfrak{M}(\mathfrak{A}))$, respectively, through the Gelfand transform.

Proposition 3.10. *Suppose some $f \in \mathscr{A}$ extends to a function F given by Theorem 3.9. Then $F : \mathfrak{M}(\mathfrak{A}) \to C(\mathfrak{M}(A))$ is continuous if and only if $\tilde{f} : \mathfrak{M}(A) \to C(\mathfrak{M}(\mathfrak{A}))$ is continuous.*

Proof. First, assume that $\tilde{f} : \mathfrak{M}(A) \to C(\mathfrak{M}(\mathfrak{A}))$. Given $\varepsilon > 0$, since $\mathfrak{M}(A)$ is compact, there exist $\phi_1, \dots, \phi_n$ in $\mathfrak{M}(A)$ such that if $\phi \in \mathfrak{M}(A)$ then $\|\widehat{\phi \circ f} - \widehat{\phi_i \circ f}\| \leq \varepsilon/3$, for some ϕ_i. Given $\psi_0 \in \mathfrak{M}(\mathfrak{A})$, define a neighbourhood V_0 of ψ_0 as follows:

$$V_0 = \{\psi \in \mathfrak{M}(\mathfrak{A}) : |\psi(\phi_i \circ f) - \psi_0(\phi_i \circ f)| < \varepsilon/3,\ 1 \leq i \leq n\}.$$

Now, let $\psi \in V_0$. For every $\phi \in \mathfrak{M}(A)$, there is ϕ_i such that

$$\begin{aligned} |\phi(F(\psi)) - \phi(F(\psi_0))| &= |\psi(\phi \circ f) - \psi_0(\phi \circ f)| \\ &\leq |\psi(\phi \circ f) - \psi(\phi_i \circ f)| + \varepsilon/3 + |\psi_0(\phi_i \circ f) - \psi_0(\phi \circ f)| \\ &\leq 2\|\widehat{\phi \circ f} - \widehat{\phi_i \circ f}\| + \varepsilon/3 \leq \varepsilon. \end{aligned}$$

Since $\phi \in \mathfrak{M}(A)$ is arbitrary, we get $\|\widehat{F(\psi)} - \widehat{F(\psi_0)}\| \leq \varepsilon$, for $\psi \in V_0$.

Conversely, suppose $F : \mathfrak{M}(\mathfrak{A}) \to C(\mathfrak{M}(A))$. Given $\varepsilon > 0$, there exist $\psi_1, \ldots, \psi_m$ in $\mathfrak{M}(\mathfrak{A})$ such that if $\psi \in \mathfrak{M}(\mathfrak{A})$ then $\|\widehat{F(\psi)} - \widehat{F(\psi_j)}\| \leq \varepsilon/3$, for some ψ_j. Take $a_j = F(\psi_j)$, for $j = 1, \ldots, m$. Given $\phi_0 \in \mathfrak{M}(A)$, take a neighbourhood U_0 of ϕ_0 as follows:

$$U_0 = \{\phi \in \mathfrak{M}(A) : |\phi(a_j) - \phi_0(a_j)| < \varepsilon/3,\ 1 \leq j \leq m\}.$$

The rest of the proof is similar to the previous part. □

Theorem 3.11. *Let $\mathscr{A}$ be an admissible Banach A-valued function algebra on X and $\mathfrak{A} = C(X) \cap \mathscr{A}$. Suppose $\|\hat{f}\| = \|f\|_X$, for all $f \in \mathfrak{A}$. Then the mapping g in Theorem 3.9 is continuous and thus every character $\psi : \mathfrak{A} \to \mathbb{C}$ lifts to some A-character $\Psi : \mathscr{A} \to A$. In particular, if $\mathfrak{A}$ is a uniform algebra, then $\mathscr{A}$ satisfies all conditions in Theorem 3.9.*

Proof. Take $\psi \in \mathfrak{M}(\mathfrak{A})$ and define $g : \mathfrak{M}(A) \to \mathbb{C}$ as in Theorem 3.9. Since $\|\hat{f}\| = \|f\|_u$, for every $f \in \mathfrak{A}$, ψ is a continuous functional on $(\mathfrak{A}, \|\cdot\|_u)$; see also [9]. By the Hahn-Banach theorem, ψ extends to a continuous linear functional $\bar{\psi}$ on $C(X)$. This, in turn, implies that g extends to a linear functional $\bar{g} : A^* \to \mathbb{C}$ defined by $\bar{g}(\phi) = \bar{\psi}(\phi \circ f)$. By Proposition 3.5, the mapping $\tilde{f} : A^* \to C(X)$, $\phi \mapsto \phi \circ f$, is w*-continuous on bounded subsets of A^*. Hence, the linear functional $\bar{g}$ is w*-continuous on bounded subsets of A^*. Since A is a Banach space, Corollary 3.11.4 in [10] shows that $\bar{g}$ is w*-continuous on A^*. □

Remark 3.12. Let $\mathscr{A}$ be an admissible A-valued uniform algebra with $\mathfrak{A} = \mathscr{A} \cap C(X)$. By Theorem 3.11, every $f \in \mathscr{A}$ extends to a function $F : \mathfrak{M}(\mathfrak{A}) \to A$. If, in addition, A is a uniform algebra so that $\|a\| = \|\hat{a}\|_{\mathfrak{M}(A)}$, by Proposition 3.10, this extension F is continuous and one can see the following maximum principle

$$\|F\|_{\mathfrak{M}(\mathfrak{A})} = \|F\|_X = \|f\|_X.$$

Example 3.13. Let K be a compact plane set. Then every $f \in P(K, A)$ has an extension $F : \hat{K} \to A$ which belongs to $P(\hat{K}, A)$, and very $\lambda \in \hat{K}$ induces an A-character $\mathcal{E}_\lambda : P(K, A) \to A$ given by $\mathcal{E}_\lambda(f) = F(\lambda)$.

Corollary 3.14. *If $\mathscr{A}$ is an admissible A-valued uniform algebra on X, then $\tilde{\mathscr{A}}$ is an admissible $\mathfrak{A}$-valued uniform algebra on $\mathfrak{M}(A)$.*

Remark 3.15. When $\mathfrak{A}$ is uniformly closed in $C(X)$, every linear functional $\psi \in \mathfrak{A}^*$ lifts to some bounded linear operator $\Psi : \mathscr{A} \to A$ with the property that $\phi(\Psi f) = \psi(\phi \circ f)$, for all $\phi \in A^*$. In fact, by the Hahn-Banach theorem, ψ extends to a linear functional $\bar{\psi} \in C(X)^*$. Then, by the Riesz representation

theorem, there is a complex Radon measure μ on X such that

$$\psi(f) = \int_X f d\mu \quad (f \in \mathfrak{A}).$$

Using [17, Theorem 3.7], one can define $\Psi(f) = \int_X f d\mu$, for every $f \in \mathscr{A}$, such that

$$\phi(\Psi(f)) = \int_X (\phi \circ f) d\mu = \psi(\phi \circ f) \quad (f \in \mathscr{A},\ \phi \in A^*).$$

References

1. M. Abel, M. Abtahi, *Description of Closed Maximal Ideals in Topological Algebras of Continuous Vector-valued Functions*, Mediterr. J. Math. **11** (2014), 1185–1193.
2. M. Abtahi and S. Farhangi, *Vector-valued spectra of Banach algebra valued continuous functions*, (in preparation)
3. F.F. Bonsall, J. Duntcan, *Complete Normed Algebras*, Springer-Verlag, Berlin, Heidelberg, New York, (1973).
4. H.X. Cao, J.H. Zhang, and Z.B. Xu, *Characterizations and extensions of Lipschitz-α operators*, Acta Math. Sin., English Series, **22** (2006), 671–678.
5. H.G. Dales, *Banach Algebras and Automatic Continuity.* London Mathematical Society Monographs, New Series 24, Oxford Science Publications, Clarendon Press, Oxford University Press, New York, (2000).
6. K. Esmaeili and H. Mahyar, *The character spaces and Śilov boundaries of vector-valued lipschitz function algebras*, Indian J. Pure Appl. Math., **45** 6 (2014), 977–988.
7. T.W. Gamelin, *Uniform Algebras*, Prentice-Hall, Inc., Englewood Cliffs, NJ., (1969).
8. A. Hausner, *Ideals in a certain Banach algebra*, Proc. Amer. Math. Soc. **8** (1957), 246–249.
9. T.G. Honary, *Relations between Banach function algebras and their uniform closures*, Proc. Amer. Math. Soc., **109** (1990), 337–342.
10. J. Horváth, *Topological vector spaces and distributions*, Vol. I, Addison-Wesley, Reading, Mass., (1966).
11. B.E. Johnson, *The uniqueness of the (complete) norm topology*, Bull. Amer. Math. Soc., **73** (1967), 537–539.
12. J.A. Johnson, *Banach Spaces of Lipschitz Functions and Vector-Valued Lipschitz Functions*, Ph.D. thesis, University of Illinois, (1969).
13. J.A. Johnson, *Banach spaces of Lipschitz functions and vector-valued Lipschitz functions*, Trans. Amer. Math. Soc., **148** (1970), 147–169.
14. E. Kaniuth, *A Course in Commutative Banach Algebras*, Graduate Texts in Mathematics, 246, Springer, (2009).
15. G.M. Leibowitz, *Lectures on Complex Function Algebras*, Scott, Foresman and Company, 1970.
16. A. Nikou and A.G. Ofarrell, *Banach algebras of vector-valued functions*, Glasgow Math. J., **56** (2014), 419–246.
17. W. Rudin, *Functional Analysis*, 2nd edn. McGraw-Hill, Inc., New York, (1991).
18. J. Tomiyama, *Tensor products of commutative Banach algebras*, Tohoku Math. J., **12** (1960), 147–154.
19. B. Yood, *Banach algebras of continuous functions*, Amer. J. Math., **73** (1951), 30–42.

20. W. Żelazko, *Banach Algebras*, Translated from the Polish by Marcin E. Kuczma. Elsevier Publishing Co., Amsterdam-London-New York; PWN–Polish Scientific Publishers, Warsaw, (1973).

Mortaza Abtahi
School of Mathematics and Computer Sciences, Damghan University, Damghan, P.O.BOX 36715-364, Iran
E-mail address: abtahi@du.ac.ir; mortaza.abtahi@gmail.com